



ACADEMY OF DENTAL LEARNING & OSHA TRAINING

Wound Healing: A New Understanding of the Drama

The Academy of Dental Learning and OSHA Training, LLC, designates this activity for 2 continuing education credits (2 CEs).

Compilation: Kristine Krafts, MD

Health Science Editor: Megan Wright, RDH, MS

Publication Date: October 2012

Updated Date: June 2020

Expiration Date: July 2023



The Academy of Dental Learning and OSHA Training, LLC is an ADA CERP Recognized Provider. ADA CERP is a service of the American Dental Association to assist dental professionals in identifying quality providers of continuing dental education. ADA CERP does not approve or endorse individual courses or instructors, nor does it imply acceptance of credit hours by boards of dentistry. Concerns or complaints about a CE provider may be directed to the provider or to the Commission for Continuing Education Provider Recognition at ADA.org/CERP.

Conflict of Interest Disclosure: ADL does not accept promotional or commercial funding in association with its courses. In order to promote quality and scientific integrity, ADL's evidence-based course content is developed independent of commercial interests. Refund Policy: If you are dissatisfied with the course for any reason, prior to taking the test and receiving your certificate, return the printed materials within 15 days of purchase and we will refund your full tuition. Shipping charges are nonrefundable.

California Registered Provider Number: RP5631

Answer Sheet: Wound Healing: A New Understanding of the Drama

- | | | | |
|----------|-----------|-----------|-----------|
| 1. _____ | 6. _____ | 11. _____ | 16. _____ |
| 2. _____ | 7. _____ | 12. _____ | 17. _____ |
| 3. _____ | 8. _____ | 13. _____ | 18. _____ |
| 4. _____ | 9. _____ | 14. _____ | 19. _____ |
| 5. _____ | 10. _____ | 15. _____ | 20. _____ |

Name: _____ Profession: _____

License State: _____ License Number: _____ Expiration Date _____

Address _____

City: _____ State: _____ Zip Code: _____

Telephone: _____ Fax: _____

E-mail: _____

If you have downloaded the course and printed the answer sheet from the Internet please enter payment information below.

Card type: _____ Card Number: _____

Exp. Date: _____ Name as it appears on card: _____

***To enter your answers online you MUST return to our website www.dentallearning.org.**

Return answer sheet:

- Via fax: 518.514.1103
- Via email: CESupport@dentallearning.com
- Postal Mail: ADL, PO Box 14585, Albany, NY 12212

*****PLEASE PRINT CLEARLY; ILLEGIBLE ANSWER SHEETS WILL NOT BE PROCESSED.**

Notes:

Course Evaluation

Please place an X in the box to rate these statements:	Poor	Fair	Good	Very Good	Excellent
The content fulfills the overall purpose of the course.					
The content fulfills each of the course objectives.					
The course subject matter is accurate.					
The material presented is understandable.					
The teaching/learning method is effective.					
The answers to the test questions are appropriately covered in the course.					
How would you rate this course overall?					
Time to complete the entire course and the test?	Hours: _____			Minutes: _____	
	Google				
	Other Search Engine				
	Friend/Coworker				
	Other				
Do you have any suggestions about how we can improve this course? If so please note them on a separate sheet of paper and send it in with your answer sheet.					
If you studied the course online, did all the links work? If not please note the page and link on a separate sheet of paper and send it in with your answer sheet so we can fix it.					

Instructions

1. Review the Objectives: Objectives provide an overview of the entire course.
2. Read the course material.
3. Complete the test:
 - a. Return to our website: www.dentallearning.org, click on Take the Exam, enter your answers, register, if you are new customer (existing customers login), pay for the course, click Grade Test. Your test will be graded immediately. If required, complete the course evaluation. Your certificate will display for you to print.
 - b. If you would rather, you may return your completed answer sheet and course evaluation to us via the options listed below.

To successfully complete the course you must score 80% or above on the test. If you do not score 80% you may retake the test one more time free of charge. If you fail a second time you must purchase a new course and test.

If you've downloaded this coursebook off the Internet you can:

- Return to our website (www.dentallearning.org) to take the test online (only if you have not purchased the coursebook separately). You will need to provide credit card information at the time you submit your test online for scoring.
- Write your answers on the one-page answer sheet included in this book, complete the credit card payment information, and return the form to the address below, fax, or email address below. Or, you may send a check or money order to the address below with your answer sheet.

Academy of Dental Learning and OSHA Training, LLC (ADL)

P.O. Box 14585

Albany, NY 12212

Fax: 518-514-1103

Email: CESupport@dentallearning.org

Answer sheets received without payment will not be processed.

We grade all tests in a timely manner; if you do not receive your certificate within five days, please email (CESupport@dentallearning.org) or call us: 518-209-9540.

There is no time limit for return of your answer sheet. Completion dates are taken from the envelope postmark or the finish date recorded in the computer when you do an online exam. Tests MUST be completed in the licensing cycle you wish to use the credits.

If you are dissatisfied with the course for any reason, prior to taking the test and receiving your certificate, return the printed materials within 15 days of purchase and we will refund your full tuition. Shipping charges are nonrefundable.

If someone else would like to use this material after you are done, he or she may register with us and take advantage of a “sharing discount”. Courses downloaded from the Internet can be shared at the same tuition rate as currently available on our website. Please call us if you need an extra answer sheet or download one from our website. There is no “sharing discount” for online exams.

The author and ADL have made every effort to include information in this course that is factual and conforms to accepted standards of care. This course is not to be used as a sole reference for treatment decisions. It is your responsibility to understand your legal obligations and license requirements when treating patients. ADL is not responsible for the misuse of information presented in this course. The material in this course cannot be reproduced or transmitted in any way without the written consent of ADL.

Table of Contents

Answer Sheet	1
Evaluation	2
Instructions	3
Table of Contents	5
Objectives	6
Introduction	6
Type of Wound Healing	6
The Actors	8
The Cells	8
The Script	10
Growth Factors	10
Cytokines	12
Receptor Mediating Signal	12
The Cell Cycle	13
The Stage	14
Extra-Cellular Matrix	14
The Drama	16
Regeneration	16
Scarring	16
First Intension vs. Second Intention Wounds	17
Act I: Inflammation	17
Act II: Proliferation	18
Act III: Maturation	20
Bad Endings in the Drama	21
Updates in Wound Healing	23
Conclusion	25
References	25
Author's Note	28
Course Exam	28

Objectives

Upon completion of this course, the student will be able to:

- List the main cells and their roles in wound healing.
- List the primary growth factors involved in tissue repair.
- Understand the messages growth factors sent to healing tissues.
- Describe the way cells are able to “hear” instructions using receptor-mediated signaling.
- Explain why the extra-cellular matrix is critical to wound healing.
- Understand the difference between tissue regeneration and scarring.
- Predict if healing will take place along the regeneration or scarring pathway.
- Know the differences between first-intention and second-intention healing.
- Understand the three phases of the wound healing.
- Describe local and systemic factors that may impede wound healing.
- Name the single most important cause of delayed wound healing.
- Identify two forms of abnormal wound healing.

Introduction

As living beings, we encounter every kind of traumatic event from paper cut to dental extraction to myocardial infarction. We must possess ways to heal damaged tissues to survive. While some animals are able to regrow complete body parts following injury (such as the earthworm who grows a new head following bisection), humans are sadly incapable of such feats. Our means of recovery following tissue damage consists largely of repair rather than pure regeneration. Thousands of times in our lives, a meticulously scripted but unseen wound healing drama is enacted, with cells serving as actors, extra-cellular matrix as the setting, and growth factors as the means of communication. This course presents the main features of this fascinating drama as it is enacted in healthy patients, and describes ways in which the drama ends badly and healing is not achieved.

Types of Wound Healing

Definitions

The term “**repair**,” when used in the context of the healing of damaged tissue, is defined as the restoration of tissue architecture and function after an injury. It encompasses two separate processes: regeneration and replacement.

Regeneration refers to a type of healing in which new growth completely restores portions of damaged tissue to their normal state. **Replacement** refers to a type of healing in which severely damaged or non-regenerable tissues are repaired by the laying down of connective tissue, a process commonly referred to as **scarring**.

A few types of tissue injury like minor paper cuts can sometimes be healed in such a way that no permanent damage remains, yet most of our tissue repair consists of both regeneration and replacement. Tissue repair may restore some of the original structures of the damaged tissue such as epithelial layers, but may also result in structural abnormalities that impair organ function. An example is the scar formed in the healing of a myocardial infarction.

Types of tissues

The healing of a wound proceeds down the pathway of regeneration or replacement – or both – depending partly on the type of tissue in which the damage occurs. Certain tissues of the body are more capable of cellular proliferation and subsequent regeneration than others.

There are three types of tissues:

- continuously dividing tissues
- quiescent tissues
- non-dividing tissues

Continuously dividing tissues (also known as labile tissues) are comprised of cells that are constantly proliferating in order to replace dead or sloughed-off cells. Examples of such tissues include epithelia (such as skin, mucosal membranes, and gastrointestinal epithelium) and hematopoietic tissues. These tissues contain pools of stem cells, which have enormous proliferative and self-renewing ability, and which give rise to more than one type of cell. Replicating asymmetrically, each stem cell gives rise to one daughter cell which differentiates and matures, and another daughter cell which remains undifferentiated and capable of beginning another self-renewing cycle.

Some tissues, known as **quiescent tissues** (or **stable tissues**) are composed of cells which normally exist in a non-dividing state but may enter the cell cycle in response to certain stimuli, such as cell injury. Tissues falling into this category include parenchymal cells of the liver, kidney, and pancreas, mesenchymal cells such as fibroblasts and smooth muscle cells, endothelial cells and lymphocytes.

It should be noted that the liver, unlike other quiescent tissues, has a relatively robust proliferative capacity. When a lobe of the liver is resected for donation, for example, the remaining liver cells proliferate at such a rate that the liver reaches a size similar to that

prior to resection. While this process is commonly described as regeneration, it is more accurately viewed as compensatory growth, since the original lobe itself does not re-grow.

A few types of tissue are composed of cells that have left the cell cycle permanently, and are therefore unable to proliferate. These **non-dividing tissues** (or **permanent tissues**) include cardiac and skeletal muscle. Tissue repair in these tissues always leaves permanent evidence of injury, such as a scar (Kumar, Abbas, Fausto, Aster, 2010).

The Actors

The Cells

The cast of characters in the tissue repair drama is large and varied. Here we list the major actors, with a focus on those that are less well-known. Central in the drama are the tissue's own lost or damaged cells, which in most cases are terminally differentiated and incapable of replication. In non-dividing tissues, such as myocardial tissue, lost cells are simply never replaced. In other tissues, however, replacement is possible. Continuously dividing tissues are particularly adept at self-renewal, undergoing innumerable cycles of cell loss and replacement during a normal human lifespan. The regenerative capacity of these tissues lies not in their parenchymal cells (which are terminally differentiated and thus unable to replicate), but in stem cells located deep within the tissue.

Stem cells

Stem cells are unique for two reasons:

- 1) The ability to self-renew
- 2) The capacity to generate more than one cell type.

Self-renewal occurs either by symmetric replication, in which a stem cell gives rise to two daughter stem cells, equally capable of self-renewal, or as asymmetric replication, in which one daughter cell remains a self-renewing stem cell, and the other daughter cell differentiates and matures.

The capacity of a stem cell to give rise to multiple lineages of cells is most striking in embryonic stem cells. These cells, which are denoted as pluripotent, are capable of generating cells from any of the tissues of the body. Adult (or somatic) stem cells are designated as multi-potent, and give rise to a more restricted array of cell types. As expected, somatic stem cells have been found in continuously dividing tissues, such as skin, gastrointestinal epithelial lining, cornea and hematopoietic tissue. However, they have also been discovered in certain quiescent tissues, such as liver, pancreas and

adipose tissue, in which they do not normally produce differentiated cells. Most surprising is the recent discovery of stem cells residing in certain parts of the central nervous system, an organ system whose tissues have long been thought to be incapable of proliferating (Conover and Notti, 2008).

Most somatic stem cells are located in niches, micro-environments within a tissue comprised of both stem cells and non-stem cells. Neighboring non-stem cells signal the stem cell to divide when necessary, a task which the stem cell generally performs very slowly. In the skin, stem cells located in a niche within the hair follicle bulge give rise to all the cells comprising the hair follicle and contribute to the production of new surface epithelial cells after wounding (Tumbar, Guasch, Greco, Blanpain, Lowry, Rendl, et al., 2004).

Stem cells of the small intestine are located within monoclonal, stem-cell derived crypts which are completely regenerated every three to five days. The bone marrow contains not only hematopoietic stem cells, which give rise to all blood cell lineages, but multipotent stromal stem cells, which travel to different regions in the body and generate chondrocytes, osteoblasts, adipocytes, myoblasts, and endothelial cells. These stromal stem cells participate in cell replenishment after tissue injury, but they do not seem to have a function in normal tissue homeostasis.

Other cells

Beyond the stem cell, three other types of cells are critical to the process of tissue repair:

- fibroblasts
- endothelial cells
- macrophages

In most wounds, complete replacement of wounded tissue to its original, unharmed state is impossible. The wound must therefore be healed using externally obtained material to reconnect the viable tissue margins. This process, discussed in detail later, involves the laying down of acellular fibrous tissue to replace the region of lost cells. The fibrous tissue is laid down by fibroblasts, which migrate to the injured area, proliferate, and secrete collagen under the influence of numerous growth factors and cytokines. In the earliest stages of wound healing, fibroblasts are few and far between, suspended together with tenuous new blood vessels in an edematous pink substance termed granulation tissue. Initially the new blood vessels are critical in the transport of nutrients and cells to the new tissue, but after a time, they recede along with the fibroblasts, leaving a collagenous scar which is remodeled and strengthened over time. Macrophages are essential directors of this drama, secreting growth factors which entice and stimulate fibroblasts, endothelial precursor cells and (in skin wounds)

keratinocytes. They also oversee the deposition and remodeling of extra-cellular matrix material.

The Script

The script of the tissue factor drama - the migration and proliferation of cells, the laying down of extra-cellular matrix, and the remodeling of collagen to form a durable scar – is carried out by a process known as receptor-mediated signal transduction. Like words spoken between people, ligands such as growth factors and cytokines float between cells, carrying directives to perform a certain action. Cells “hear” these words when the ligands bind to cell-surface receptors, which bring the message into the cell, resulting in a new action, such as migration, proliferation, or secretion of a substance. Here we will discuss the words spoken between the actors, the way the actors hear these words, and the manner in which the message gets to the cell nucleus in order to effect change.

Growth factors and cytokines

Growth factors are specialized polypeptide molecules which bind to receptors on target cells and deliver messages regarding migration, proliferation, differentiation, survival, and secretion. The list of growth factors and their attendant functions is so long that it would tax even the most capable memorizer. Herein we limit our discussion to the primary growth factors associated with each stage of tissue repair (Table 1).

	EGF	FGF	KGF	PDGF	TGFa	TGFb	TNF	VEGF
Fibroblast migration		X		X		X		
Fibroblast proliferation	X	X		X	X		X	
Monocyte migration		X		X		X	X	
Macrophage activation							X	
Epithelial migration	X	X	X		X			
Epithelial proliferation	X	X	X		X			
Angiogenesis		X		X	X			X
Collagen synthesis				X		X		
Collagenase synthesis	X	X		X			X	X
Wound contraction		X		X				

EGF, epidermal growth factor; FGF, fibroblast growth factor; KGF, keratinocyte growth factor; PDGF, platelet-derived growth factor; TGF, transforming growth factor; TNF, tumor necrosis factor; VEGF, vascular endothelial growth factor.

Transforming Growth Factor Beta (TGF-β)

One of the most critical growth factors in tissue repair is transforming growth factor beta (TGF-β). This growth factor belongs to a superfamily which also includes a number of other factors with wide-ranging functions, such as bone morphogenetic proteins, activins, inhibins, and Müllerian inhibiting substance (Bessa, Casal, Reis, Bone, 2008).

Made by platelets, endothelial cells, lymphocytes and macrophages, TGF- β binds to cell-surface receptors which have serine/threonine kinase activity, triggering phosphorylation of cytoplasmic transcription factors called Smads, so-named for the two Drosophila proteins of which they are homologues: the mothers against decapentaplegic (MAD) and Caenorhabditis elegans proteins (Itoh, Asao, Sugamura, et al., 2001). Smads then enter the nucleus and affect gene transcription.

Although TGF- β has many functions, its most important role in tissue repair is to promote fibrosis, a feat it accomplishes by:

- Attracting fibroblasts and stimulating them to proliferate
- Triggering fibroblasts to secrete collagen
- Inhibiting extra-cellular matrix degradation by metalloproteinases.

Not surprisingly, TGF- β is a key factor in conditions involving pathologic fibrosis, such as pulmonary fibrosis, systemic sclerosis, and Marfan syndrome (Horan, Wood, Ona, Li, Lukashev, Weinreb, et al., 2001). In addition to its fibrogenic effects, TGF- β also inhibits epithelial cell growth, diminishes inflammation, and promotes invasion and metastasis in tumors (Attisano, Wrana, 2002).

Platelet-Derived Growth Factor (PDGF)

Another essential, multitasking growth factor is platelet-derived growth factor (PDGF), so named because it is stored in platelet granules and released upon platelet activation. Made by a number of cells including macrophages, endothelial cells and smooth muscle cells, PDGF is involved in nearly every aspect of tissue repair. It calls neutrophils, macrophages and fibroblasts to the wound area, and subsequently stimulates and activates them. It also induces angiogenesis, triggers production of matrix metalloproteinases (MMPs), fibronectin and hyaluronic acid, and aids in wound contraction (Andrae, Gallini, Betsholtz, 2008).

Fibroblast growth factor (FGF)

Fibroblast growth factor (FGF) calls virtually all of the major players (macrophages, fibroblasts, and endothelial cells) to the scene of the wound. It initiates the migration of epithelial cells in from the margins of the wound, mediates wound contraction, and stimulates angiogenesis (Brakenhielm, Pawliuk, Wariaro, Post, Wahlberg, et al., 2003). Contrary to what one might infer from its name, however, FGF does not stimulate collagen synthesis. The more appropriately-named vascular endothelial growth factor (VEGF) acts primarily on endothelial cells. It is a potent inducer of blood vessel formation in tissue repair as well as in early fetal development (Holmes, Roberts, Thomas, Cross, 2007). Overexpressed in certain tumors, particularly renal cell carcinoma, VEGF is a target for the development of new chemotherapy drugs (Escudier

B., 2007). Other growth factors include epidermal growth factor (EGF) and hepatocyte growth factor (HGF). Both stimulate epithelial cell and hepatocyte proliferation and enhance epithelial cell migration. Once known as the scatter factor, HGF also promotes cell scattering during embryonic development (Schmidt, Bladt, Goedecke, Brinkmann, Zschiesche, Sharpe, et al., 1995.)

Cytokines

In addition to growth factors, cytokines also carry important messages between cells during tissue repair. Small proteins secreted by cells of the immune system, cytokines are perhaps best known for their role as immunomodulators within both the cellular and humoral arms of the adaptive immune system. Three cytokines in particular are involved in tissue repair: tumor necrosis factor (TNF) and interleukin-1 (IL-1) participate in the wound healing stage of tissue repair, and TNF and interleukin-6 (IL-6) are involved in liver regeneration (Fedyk, Jones, Critchley, Phipps, Blieden, Springer, 2001).

Receptor-mediated signaling

For humans, words must be seen or heard to have meaning. So it is with the messages between cells in the tissue repair drama: the growth factor or cytokine message must have a way to get into the target cell for action to take place. In cells, this process has been given the somewhat cumbersome term *receptor-mediated signal transduction*, and it involves binding of a ligand (a growth receptor or cytokine) to a receptor molecule on a target cell. This binding triggers events which transduce the signal from the outside of the cell to the inside. The end result is a change in gene expression, for example, transcription of genes that normally may be silent, such as those that control cell cycle entry.

Receptor-mediated signaling often occurs between cells adjacent to one another (paracrine signaling). For example, macrophages produce growth factors which bind to receptors on adjacent fibroblasts. However, signaling can also occur between cells located at some distance from each other (endocrine signaling), as is often the case with cytokines and their distant targets. Signaling may even occur within the same cell (autocrine signaling), as when a tumor cell overproduces both a growth factor and its receptor, which interact and lead to unbridled cell growth.

Whatever the mode of signaling may be, there are four main types of receptors: those with intrinsic tyrosine kinase activity, those without intrinsic tyrosine kinase activity, G protein-coupled receptors, and steroid hormone receptors. Receptors with intrinsic tyrosine kinase activity are common targets for growth factors. When a growth factor binds to this type of receptor, the receptor dimerizes, causing phosphorylation of tyrosine residues and activation of the receptor. The activated receptor in turn phosphorylates other molecules, inducing them to carry the message (or "signal") into

the nucleus, resulting in a change of gene expression. Examples of actions mediated by this type of receptor include: production of growth factors, production of growth factor receptors, production of proteins that control entry of the cell into the cell cycle, and activation of cell proliferation and survival (through inhibition of apoptosis) (Robinson, Wu, Lin, 2001).

Receptors lacking intrinsic tyrosine kinase activity use a similar phosphorylation-activation system to transmit messages into the cell; however, since they lack phosphorylating capability, they must recruit a separate kinase system, called the JAK-STAT pathway, to complete the job. The receptor transmits the signal from the ligand (often a cytokine) on the surface of the cell to a Janus kinase (JAK) protein inside the cell. Once activated, the JAK protein in turn phosphorylates and activates cytoplasmic transcription factors called STATs (signal transducer and activator of transcription). As their name implies, STATs transduce the signal from the JAK protein to the nucleus of the cell, where they activate the transcription of certain genes (Rawlings, Rosler, Harrison, 2004).

The largest family of signal transduction cell membrane receptors is comprised of G protein-coupled receptors. Many different types of ligands bind to this type of receptor, such as chemokines, vasopressin, serotonin, histamine, epinephrine, norepinephrine, calcitonin, glucagon, parathyroid hormone, corticotropin and rhodopsin.¹⁶ When the receptor receives a signal from a ligand, its seven transmembrane alpha helices change conformation, allowing it to interact with a G protein. Once activated, the G proteins transmits the signal to a second messenger molecule, such as 3', 5' cyclic adenosine monophosphate (cAMP), which carries the signal to the nucleus of the cell. Such cAMP-mediated pathways are involved in vision and olfactory sensing. Steroid hormone receptors are unique in that they are located inside the cell rather than on the cell membrane. Their ligands, which include steroid hormones, thyroid hormone, vitamin D and retinoids, must diffuse through the cell membrane in order to bind to the receptor, which binds to DNA and alters gene expression.

The Cell Cycle

A crucial element in the tissue repair script is cell proliferation. In order to heal after injury - whether by regeneration or scarring - cells must enter and progress through the cell cycle, a tightly-regulated process that consists of two main activities: DNA replication and mitosis. Continuously proliferating cells are always moving through the cell cycle, whereas quiescent cells must be called into the cell cycle by growth factors or cytokines (via receptor-mediated signal transduction) or by ECM components (via integrins).

The cycle consists of four consecutive phases: the G1 (presynthetic) phase, the S (DNA synthesis) phase, the G2 (premitotic) phase, and the M (mitotic) phase. Cells may begin their journey through the cycle from G1 or G0 (a resting phase outside the cell cycle). In order to move from G0 to G1, cells must activate numerous previously-silent genes including proto-oncogenes, genes required for ribosome synthesis, and genes required for protein translation. To move into the S phase and begin the process of replication, cells must pass a checkpoint at the end of G1. Several such checkpoints built into the cycle operate as quality-control monitors, validating that the cell has accurately completed one phase of the cell cycle before allowing the cell to progress to the next phase. The G1 checkpoint assesses the integrity of DNA before allowing cells to pass through to S phase. It also serves as a decision point, determining whether the cell should divide immediately, divide at a later point, or enter a resting stage. Another checkpoint at the end of G2 evaluates DNA after replication to see if the cell can safely enter mitosis. A separate metaphase checkpoint monitors the alignment of chromosomes at the mitotic plate, and when alignment is achieved, allows chromosomes to separate into their sister chromatids. After splitting into two daughter cells, the cell again enters G1.

Because the correct functioning of the cell cycle is crucial to the integrity of the cell, there are multiple controls built into the system, including activators, inhibitors, and sensors that control checkpoints in the cycle. Proteins called cyclins, together with associated cyclin-dependent protein kinases (CDKs), drive the cell cycle by phosphorylating proteins that are essential for cell cycle transitions.¹⁷ One such protein is the retinoblastoma (RB) protein, which in its normal, unphosphorylated state stalls the cell at the G1/S transition by binding and inactivating the transcription factor E2F.¹⁸ When RB is phosphorylated by cyclins, it releases (and activates) E2F, allowing transcription of genes necessary for cell cycle progression. Cyclin proteins are themselves regulated by CDK inhibitors.

The Stage

Functions of the Extracellular Matrix

The stage upon which the drama unfolds is a special type of tissue known as extracellular matrix (ECM). ECM exists in two forms: interstitial matrix (a gelatinous, amorphous intercellular material) and basement membrane (a thin, highly-organized, plate-like layer upon which epithelial cells rest). Although it may appear inert and static, ECM has a long list of responsibilities and functions. Beyond its obvious physical characteristics - imparting resilience to soft tissues and firmness to bone - it also stores and presents growth factors, acts as a scaffold to which migrating cells can adhere and

establish polarity, and facilitates cell growth both in physiologic and tissue repair settings. ECM is the ever- changing background for regeneration and wound healing, but it also accompanies other processes such as morphogenesis, chronic fibrotic processes, and tumor invasion/metastasis.

The many constituents of the ECM can be grouped into three types of molecules, according to their main functions: structural molecules (such as collagen and elastin), adhesive molecules (such as integrins, fibronectin and laminin), and resilient components (such as proteoglycans and hyaluronan). The interstitial matrix is composed primarily of collagen, elastin, fibronectin, proteoglycans, and hyaluronan, whereas the basement membrane is composed largely of nonfibrillar collagen, laminin, and proteoglycans (Kumar, Abbas, Fausto, Aster, 2010). A short description of each major ECM component follows.

Components of the Extra-cellular Matrix

Collagen, the most common protein in the animal world, consists of three chains which combine to form a triple helix trimer. Although 27 types of collagen have been identified, less than half of these have major roles in the human body. Fibrillar collagens (types I, II, III, V, and IX) are found in many types of hard and soft tissues. Type IV collagen, which is arranged in long sheets rather than fibrils, is a primary component of basement membranes. Collagen imparts strength to tissues, but the ability to stretch and snap back into shape is provided by elastic fibers. Elastic fibers are composed of a central core of elastin surrounded by fibrillin, a glycoprotein which controls the availability of TGF- β within the ECM.

Adhesive molecules, such as integrins, fibronectin and laminin, provide connections between cells, or between cells and ECM components. Integrins are transmembrane glycoproteins with an extracellular domain that attaches to ECM components, such as laminin and fibronectin, and an intracellular domain that links to cytoskeletal complexes (Hynes. Integrins, 2002). They are operative in the earliest stages of wound healing, helping leukocytes adhere to vascular endothelium in preparation for their transmigration through the vessel. Fibronectin helps stabilize the initial blood clot filling the wound by binding to fibrin; it also provides a framework for building the collagen matrix during wound healing. Laminin provides a connection between cells and the ECM; it also binds with type IV collagen to form the basement membrane.

The resilient nature of the ECM is provided by proteoglycans and hyaluronan. Both of these substances are composed of glycosaminoglycans (GAGs), long repeating polymers of disaccharides. Proteoglycans, which consist of GAGs linked to a protein core, have many different roles: regulation of ECM permeability, mediation of inflammatory and immune responses, and control of cell growth. Hyaluronan,

comprised of many repeats of a single disaccharide, binds huge amounts of water and provides strength and turgor to connective tissue.

The Drama

Regeneration

True regeneration - in which new cells replace damaged or dead cells so that the tissue is restored to its original state - occurs infrequently in humans. Two conditions must be met for a tissue to undergo pure regeneration (without scarring):

- 1) the injured cells must be capable of proliferation (as is the case in continuously dividing and quiescent tissues), and
- 2) the underlying stromal framework must be intact.

Regeneration occurs in a physiologic manner in continuously dividing tissues. Sloughed-off gastrointestinal epithelial cells, for example, are replaced by new cells arising from stem cells in the intestinal crypts. Pathologic examples of pure regeneration, however, are few and far between. One example occurs when the liver undergoes acute, toxic injury from acetaminophen overdose. In this type of injury, hepatocytes are destroyed, but the underlying stromal framework remains intact, allowing for the possibility of complete regeneration of the injured tissue (Apte, Singh, Zeng, Cieply, Virji, Wu, et al., 2009).

In a process akin to regeneration, some organs are able to grow in size in response to resection. If a lobe of the liver is resected, the remaining liver will grow larger in response. This process, though often termed “regeneration,” is not true regeneration, but compensatory growth. True regeneration of lost organs is, at this time, a process relegated to certain animal species and science fiction movies – but perhaps in the not-so-distant future stem cells will be used for this purpose (Apte, Singh, Zeng, Cieply, Virji, Wu, et al., 2009).

Scarring

If an injury damages only parenchymal cells in a continuously-dividing or quiescent tissue, repair by regeneration is possible. However, if an injury is so severe as to damage not only the parenchymal cells but the underlying stromal framework of the tissue (as occurs in most injuries), or if the injury occurs in non-dividing tissues, regeneration is impossible. In these instances, the tissue is repaired by the deposition of fibrous tissue – a scar – in the defect created by the wound. Regeneration involves restitution of tissue components; repair involves “patching” rather than restoring. The amount of regeneration vs. repair that occurs depends on the proliferative capacity of the cells, the integrity of the stromal framework, and the duration of the injury and

inflammatory response (Kumar, Abbas, Fausto, Aster, 2010).

Repair by scarring involves the influx of debris-removing inflammatory cells, formation of granulation tissue (a substance consisting of fibroblasts and delicate capillaries in a loose extra-cellular matrix), and conversion of said granulation tissue into fibrous tissue which is remodeled over time to form a scar. There are five major components in this process: inflammation, new vessel formation, fibroblast proliferation, collagen synthesis, and scar remodeling.

First-Intention vs. Second-Intention Wounds

Though the general principles are the same no matter what type of wound is being healed, the extent of granulation tissue, inflammation and scarring vary depending on the size and type of wound. In the case of skin wounds, there are two types of wound healing: first-intention healing and second-intention healing (Broughton, Janis, Attinger, 2006).

Wounds healing by first intention are relatively small, with limited epithelial and connective tissue damage, such as paper cuts, well-approximated surgical incisions and superficial stab wounds. Healing in this type of wound is generally rapid, since the area of the defect is relatively small. Inflammation and granulation tissue are present but not abundant, and scarring is minimal.

In contrast, wounds healing by second intention are larger wounds with margins that are not easily approximated, such as extraction sockets, burns, and large excisional skin wounds. This type of wound necessarily heals more slowly, as the defect to be repaired is larger. The abundance of fibrin, necrotic debris, and exudate necessitates a more intense inflammatory response, and consequently the chance of inflammation-mediated injury is greater than it is in first-intention healing. Granulation tissue is abundant, and wound contraction and scarring is maximal. Despite these differences between first and second intention healing, the underlying drama is very similar. We will present the drama as it occurs in first intention wounds, noting second intention differences where necessary.

The Drama in Three Acts

Act I: Inflammation

Immediately after the wound is inflicted, the most urgent task is not to repair the damaged tissue, but to stop the flow of blood from the wound. Within seconds of the injury, exposed collagen activates coagulation in the region of the injury. Platelets form an initial plug, coagulation cascade factors interact to form fibrin (which seals and solidifies the plug), and a blood clot soon appears at the skin surface. Not only does the

clot stop the bleeding, but it serves as a scaffold for incoming cells and a repository of cytokines and growth factors (Broughton, Janis, Attinger, 2006) (Witte, Barbul, 1997).

At the skin surface, all appears static and calm as the clot dries and forms a scab. Underneath the surface, however, a flurry of activity begins. An army of neutrophils rushes to the scene through dilated local blood vessels. Within 24 hours, the neutrophils appear at the edges of the wound incision, using the fibrin scaffolding of the clot to invade the region of destruction, their primary tasks being to break down debris and kill bacteria.

Epithelial cells, meanwhile, slowly begin to crawl from the basal layers of the epidermis into the wounded region, depositing basement membrane components as they migrate. They are called to begin this process by growth factors secreted by platelets, macrophages and fibroblasts (Smola, Thiekötter, Fusenig, 1993) (Xia, Zhao, Marcus, Jiminez, Ruben, Moore, et al., 1999). Fusing in the midline underneath the inert scab, these epithelial cell pioneers form the first thin, continuous layer of epithelium upon which the rest of the new epithelium will be built (Lawrence WT, Diegelmann, 1996).

Act II: Proliferation

Following the initial flurry of clotting and neutrophil activity, the work of cell proliferation begins in earnest. The characteristic feature of this act in the drama is the formation of a substance called granulation tissue, a pink, soft material so-named for its irregular, grainy appearance. Consisting of new blood vessels and fibroblasts in an extra-cellular matrix, granulation tissue is laid down over a period of a few days, beginning with a few small, tenuous blood vessels and scattered fibroblasts. Initially, the new vessels are leaky, and the granulation tissue has an edematous, loose appearance (Figure 1). As the vessels grow stronger, and fibroblasts begin to lay down collagen, it takes on a more substantial appearance, reaching its maximum vascularity by day five (Kumar, Abbas, Fausto, Aster, 2010).




Figure 1. Skin ulcer showing early-stage granulation tissue with numerous blood vessels and scattered fibroblasts in a loose extracellular matrix. Numerous neutrophils are present within the granulation tissue.

Macrophages - large, multitasking cells derived from monocytes - now replace the neutrophils that were so numerous in the immediate aftermath of the wounding. As their name suggests, a primary function of macrophages is the ingestion of unwanted materials: bacteria, cell remnants, debris, fibrin, and foreign material. But macrophages also serve as directors for many other parts of the drama. They call in and stimulate fibroblasts and keratinocytes, stimulate the formation of new blood vessels, and direct the laying down and remodeling of extra-cellular matrix (Broughton, Janis, Attinger, 2006).

The formation of new blood vessels (angiogenesis) begins within the first few days following injury (WernerS, Grose, 2003). The process is unexpectedly aided by a small number of bone-marrow-derived endothelial precursor cells which home to the wound by mysterious mechanisms. Their contribution to the overall angiogenesis effort, however, is minimal (Bluff, Ferguson, O’Kane, Ireland, 2007). Formation of the correct blood vessel structure is aided by integrins on the surface of endothelial cells; surrounding perithelial cells are recruited with the help of angiopoietin.

The end point toward which the entire drama is directed is the formation of a scar, a strong collagen filler that bridges the gap left by tissue destruction, restoring strength and integrity to the tissue. This process begins with the entrance of the fibroblast between days one and three. At first, the collagen secreted by fibroblasts is arrayed vertically at the margins of the incision. Later, the it combines with fibrin and plasma fibronectin, forming a provisional scaffolding for further fibroblast and endothelial cell ingrowth (Figure 2) (Pierce, Mustoe, Altrrock, Deuel, Thomason, 2004).




Figure 2. Early granulation tissue with blood vessels in a loose matrix of collagen and fibrin.

By day three to day five, ECM formation is well underway, with granulation tissue serving as a substrate (Figure 3).




Figure 3. Skin ulcer showing later-stage granulation tissue with blood vessels and early collagen deposition.

Shards of elastic tissue form within the wound site. Plump fibroblasts mature into thin, spindle-shaped fibrocytes, which become entrapped in the dense, fibrillar collagen they have secreted (Figure 4). Collagen deposition is a balance between synthesis and degradation. In these early stages, the balance tips toward collagen synthesis. The process of ECM deposition ends within two to three weeks.





Figure 4. Early scar formation showing fibrillar collagen, plump fibroblasts and spindle-shaped fibrocytes.

Meanwhile, the epithelium quietly continues to proliferate. It increases in thickness, eventually regaining its full, stratified architecture and keratinized surface some time between one and three weeks (or later in second-intention healing) (Werner, Grose, 2003).

Act III: Maturation

During the second week following the incision, leukocytes gradually abandon the wound area and vessels recede (Figure 5). Type I collagen synthesis increases, due to both an increased number of fibroblasts and an increased rate of synthesis per fibroblast.³⁰ Where there was once granulation tissue, there is now a preliminary scar, pale and firm, composed of dense collagen, fibroblasts, and shards of elastic tissue. In another two weeks, even the fibroblasts begin to disappear, and eventually an acellular scar bridges the region of the wound, covered by intact, fully-developed epidermis.

Figure 5. Wide excision of skin.



Over the next several weeks, the composition of the extra-cellular matrix changes, remodeling the newly-formed scar such that it attains maximal strength at an appropriate size (Pilcher , Wang, Qin, Parks, Senior, Welgus, 2006). Ultimately, strength is achieved by increased collagen synthesis, followed by post-synthetic modifications of collagen, including cross- linking and increased fiber size. Early in wound healing, the collagen is relatively thin and oriented parallel to the skin; over time, the initial collagen fibrils are resorbed and replaced with thicker fibrils aligned with stress lines (Broughton, Janis, Attinger , 2006). Wound strength increases rapidly during the first month, then slows, reaching a plateau of 70-80% of the original tensile strength by the end of the third month. Although full-thickness epithelial regeneration is possible, any skin appendages lost during the initial injury will not be reformed.

Given their easily-approximated margins, wounds healing by first intention generally have minimal scarring. Wounds healing by second intention, however, are larger, with more epithelial destruction, necessitating more collagen deposition to close the wound. The process of wound contraction eases the fibroblasts' burden a bit. Specialized contractile, smooth-muscle-like cells called myofibroblasts appear within the wound and pull the margins of the wound toward each other. So efficient are these myofibroblasts that during the process of contraction, large wounds may decrease in size by up to 80% (Lorentz , Longaker, 2008).

Bad Endings in the Drama

Systemic factors affecting wound healing

Several systemic factors may adversely affect the performance of the tissue repair drama. Patients with diabetes mellitus tend to have less granulation tissue formation, less collagen deposition, and defects in collagen maturation, leading to slow and ineffective wound maturation.³³ While this predisposition has been attributed to the

microangiopathy inherent in diabetes, recent research indicates that other factors, such as collagen glycosylation and pericapillary albumin deposition, may play a role (Broughton, Janis, Attinger, 2006) (Bucalo, Eaglstein, Falanga, 2002). Poor nutrition also retards the healing process. Vitamin C, in particular, is necessary for collagen synthesis. Deprivation of this vitamin results in failure of activation of proline and lysine hydroxylases and formation of unhydroxylated procollagen peptides, resulting in an unstable, poorly cross-linked collagen helix. Conditions in which blood flow is compromised - such as cardiac insufficiency or arteriosclerosis - delay the delivery of necessary cells and factors to the scene of the wound. Certain drugs may also slow wound healing; glucocorticoids, for example, inhibit inflammation and collagen formation.

Local factors affecting wound healing


Local factors can also greatly affect the quality of wound healing. The size of the wound is important (small injuries heal faster than large ones) as is the location (wounds in richly-vascularized areas of the body, such as the face, heal relatively quickly). The presence of foreign bodies may prolong healing, as may motion or pressure at the wound site. However, the single most important cause of delayed healing of wounds is infection (Kumar, Abbas, Fausto, Aster, 2010). If any beta-hemolytic *Streptococcus* organisms are present, or if there are over 10⁵ organisms of any bacterial species per gram of tissue, the wound will not heal (Robson, 1997). More common in second-intention healing, infection prolongs the inflammatory phase of wound healing, impeding epithelialization, collagen deposition, and wound contraction. Bacterial endotoxins stimulate collagenase secretion, leading to degradation of not only the forming scar, but of surrounding normal tissue (Broughton, Janis, Attinger, 2006).

Abnormally Exuberant Wound Healing

On the other hand, sometimes acts in the drama are played so well - so exuberantly - that the end result is an abnormally-healed wound. Occasionally, the balance of collagen formation and degradation tips in favor of formation, leading to hypertrophic scars or keloids. Hypertrophic scars are raised scars that remain confined to the region of the wound. They generally occur within four weeks of injury (often a severe traumatic or thermal injury to the dermis), and may regress over time (Broughton, Janis, Attinger, 2006) (Dabiri, Tumbarello, Turner, Van De Water, 2008). An aberrant autocrine loop in which myofibroblasts produce excessive TGF- β , and hence collagen, appears to contribute to their formation (Dabiri, Tumbarello, Turner, Van De Water, 2008). Keloids are scars that have overgrown the boundaries of the initial incision, presenting as nodules or masses of fibrous tissue (Figure 6). While keloids generally appear within one year of the inciting injury, some may begin growing years later (Broughton, Janis, Attinger, 2006). For unknown reasons, keloids occur much more frequently in patients

with darkly-pigmented skin; certain people seem to have a predisposition towards their formation.


Figure 6:



Keloid showing dense, haphazardly-arranged collagen bundles.

Formation of granulation tissue may likewise occur in an excessive fashion. In a lesion known as proud flesh, the process of wound healing is interrupted by masses of granulation tissue protruding above the skin surface, preventing re-epithelialization and appropriate scar formation (Figure 7).

Figure 7:



Proud flesh with exuberant granulation tissue.

Updates in Wound Healing

Stem Cell Research

There are two therapies, currently, that utilize stem cell sheets derived from the basal layer of the skin for skin grafts, according to the Canadian Stem Cell Foundation. Primarily used to treat large burns, both treatments have been approved in the United States for clinical use.

Senior citizens with cutaneous injuries may witness a slower progression in the restoration of their wounds due to reduced skin elasticity, slower collagen replacement and age-related diseases, especially those that affect blood flow (Advanced Tissue

2017).

As a result, doctors frequently search for methods to speed up the healing process, and stem cell therapy may do the trick. These organisms stimulate the formation of blood vessels and naturally adjust the amount of inflammation, according to a study published in the journal *Gerontology*. While the publication cites the need for additional research into the identification and isolation of the most therapeutic cells, as well as an effective delivery method for cell protection for this particular purpose, the results of this analysis show promise (Advanced Tissue 2017).

As scientists keep studying these miraculous cells, more advancements will be made in wound healing, and beyond!

Genetic Research

On March 2, 2016, the Biophysical Society publishes that:

“Researchers at Ohio State University have pinpointed a human gene product that helps to regulate wound healing and may control scarring in people recovering from severe injuries and damage to certain internal organs.

The protein, MG53, travels throughout the bloodstream and helps the body fix injuries to the skin, heart, lungs, kidneys and other organs without causing scars. It's a discovery that could help heal open wounds, decrease recovery time after surgery and reduce the spread of infections.

"A massive scar on your skin may look bad, but imagine you have a heart attack and get a scar on your heart--that could be lethal," says Jianjie Ma, a physiologist at Ohio State and co-author of the presentation.

All animals carry this gene, he said, and it's almost identical no matter which species. MG53 fixes the cell and tissue damage that occurs during everyday living. Even simple actions, like walking or typing, will cause injuries to the body. Usually this isn't a problem because MG53 can make repairs before there's any serious harm.

Ma and his team genetically engineered mice without the gene that makes MG53 to see what would happen without its healing capabilities. The experiments showed that the mice lacking MG53 had difficulty recovering from injury, because of their compromised repair capacity; their heart would not function well under stress conditions.

MG53 works in tandem with another protein called TGF Beta, a type of "cytokine" protein that also heals wounds, but the healing process happens so quickly that it

causes scars. If you have more TGF Beta in your bloodstream than MG53, you scar easily.

Ma's goal is to develop a therapy that will inhibit TGF Beta and promote MG53. Medical professionals can use the therapy during procedures to promote quick, scarless healing. His next step is to identify a small compound that can do this and eventually test whether it has the desired effect in human trials.”

Conclusion

Wound healing is a carefully-scripted drama performed countless times throughout a human lifespan. Though occasionally the drama does not go as planned, in most cases damage is contained, dead cells and tissue are removed, and a scar restores integrity to the injured tissue – a triumphant performance indeed.

References

Advanced Tissue. How Stem Cells Benefit Wound Healing. April 20, 2017
<https://www.advancedtissue.com/cutting-edge-techniques-for-chronic-wound-healing/>
Accessed April 2017.

Andrae J, Gallini R, Betsholtz C. Role of platelet-derived growth factors in physiology and medicine. *Genes Dev* 2008; 22(10):1276-1312.

Apte U, Singh S, Zeng G, Cieply B, Virji MA, Wu T, et al. Beta-catenin activation promotes liver regeneration after acetaminophen-induced injury. *Am J Pathol* 2009; 175(3):1056-65.

Attisano L, Wrana JL. Signal transduction by the TGF-beta superfamily. *Science* 2002; 296(5573):1646-1647.

Bessa PC, Casal M, Reis RL. Bone morphogenetic proteins in tissue engineering: the road from laboratory to the clinic. Part 1 – basic concepts. *J Tiss Eng Regen Med* 2008; 2(1):1-13.

Biophysical Society. "Gene identified that helps wound healing: New research on gene that regulates healing and may control scarring,." *ScienceDaily*, 2 March 2016.
www.sciencedaily.com/releases/2016/03/160302135149.htm Accessed April 2017.

Bluff J, Ferguson M, O’Kane S, Ireland G. Bone marrow-derived endothelial progenitor cells do not contribute significantly to new vessels during incisional wound healing. *Exp Hematol* 2007; 35(3):500-506.

Broughton G, Janis JE, Attinger CE. Wound healing: An overview. *Plast Reconstr Surg* 2006(suppl); 117:1-32.

Bucalo B, Eaglstein WH, Falanga V. Inhibition of cell proliferation by chronic wound fluid. *Wound Rep Regen* 2002; 1(3):181-186.

Cao R, Brakenhielm E, Pawliuk R, Wariaro D, Post MJ, Wahlberg E, et al. Angiogenic synergism, vascular stability and improvement of hind-limb ischemia by a combination of PDGF- BB and FGF-2. *Nat Med* 2003; 9(5):604-613.

Cherezov V, Rosenbaum DM, Hanson MA, Rasmussen SG, Thian FS, Kobilka TS, et al. High-resolution crystal structure of an engineered human b2-adrenergic G protein-coupled receptor. *Science* 2007; 318(5854):1258-1265.

Chow LW, Loo WT, Yuen KY, Cheng C. The study of cytokine dynamics at the operation site after mastectomy.

Conover JC, Notti RQ. The neural stem cell niche. *Cell Tissue Res* 2008; 331:211-224.

Dabiri G, Tumbarello DA, Turner CE, Van De Water L. Hic-5 promotes the hypertrophic scar myofibroblast phenotype by regulating the TGF-b-1 autocrine loop. *J Invest Dermatol* 2008; 128:2518-2525.

Diegelmann R. Analysis of collagen synthesis. *Methods Mol Med* 2003; 78:349-358.

Escudier B. Anti-VEGF therapy for renal cell carcinoma. *Clin Adv Hematol Oncol* 2007; 5(7):530-531.

Fedyk ER, Jones D, Critchley HO, Phipps RP, Blieden TM, Springer TA. Expression of stromal-derived factor-1 is decreased by IL-1 and TNF and in dermal wound healing. *J Immunol* 2001; 166:5749-5754.

Holmes K, Roberts OL, Thomas AM, Cross MJ. Vascular endothelial growth factor receptor-2: Structure, function, intracellular signaling and therapeutic inhibition. *Cell Signal* 2007; 19(10):2003-2012.

Horan GS, Wood S, Ona V, Li DJ, Lukashev ME, Weinreb PH, et al. Partial inhibition of integrin avb prevents pulmonary fibrosis without exacerbating inflammation. *Am J Resp Crit Care Med* 2008; 177:56-65.

Hynes RO. Integrins: bidirectional, allosteric signaling machines. *Cell* 2002 110(6):673-687.

Itoh F, Asao H, Sugamura K, Heldin CH, ten Dijke P, Itoh S. Promoting bone morphogenetic protein signaling through negative regulation of inhibitory Smads. *EMBO J* 2001; 20(15):4132-4142.

Kumar V, Abbas AK, Fausto N, Aster JC. Tissue Renewal, Regeneration and Repair. In:

Robbins and Cotran Pathologic Basis of Disease. Eighth edition. Philadelphia: Elsevier, 2010:79-110.

Lawrence WT, Diegelmann RF. Growth factors in wound healing. *Clin Dermatol* 1994; 12(1):157-169.

Lorentz HP, Longaker MT. Wounds: Biology, pathology and management. In: Norton JA, Barie PS, Bollinger RR, Chang AE, Lowry S, Mulvihill SJ, et al., eds. *Surgery: Basic science and clinical evidence*. 2nd edition. New York: Springer, 2008:191-208.

Münger K, Howley PM. Human papillomavirus immortalization and transformation functions. *Virus Res* 2002 89(2):213-228.

Nigg EA. Cyclin-dependent protein kinases: key regulators of the eukaryotic cell cycle. *Bioessays* 1995; 17(6):471-480.

Pierce GF, Mustoe TA, Altrock BW, Deuel TF, Thomason A. Role of platelet-derived growth factor in wound healing. *J Cell Biochem* 2004; 45(4):319-326.

Pilcher BK, Wang M, Qin X, Parks WC, Senior RM, Welgus HG. Role of matrix metalloproteinases and their inhibition in cutaneous wound healing and allergic contact hypersensitivity. *Ann NY Acad Sci* 2006; 878:12-24.

Rafii S, Lyden D. Therapeutic stem and progenitor cell transplantation for organ vascularization and regeneration. *Nat Med* 2003; 9:702-712.

Rawlings JS, Rosler KM, Harrison DA. The JAK/STAT signaling pathway. *J Cell Science* 2004; 117:1281-1283.

Robinson DR, Wu YM, Lin SF. The protein tyrosine kinase family of the human genome. *Oncogene* 2000; 19(49):5548-5557.

Robson MC. Wound infection: A failure of wound healing caused by an imbalance of bacteria. *Surg Clin North Am* 1997; 77(3):637-650.

Schmidt C, Bladt F, Goedecke S, Brinkmann V, Zschesche W, Sharpe M, et al. Scatter factor/hepatocytes growth factor is essential for liver development. *Nature* 1995; 373:699-702.

Smola H, Thiekötter, Fusenig NE. Mutual induction of growth factor gene expression by epidermal-dermal interaction. *J Cell Biol* 1993; 122(2):417-429.

Tumbar T, Guasch G, Greco V, Blanpain C, Lowry W, Rendl M, et al. Defining the epithelial stem cell niche in skin. *Science* 2004; 303(5656):359-363.

Werner S, Grose R. Regulation of wound healing by growth factors and cytokines.

Physiol Rev 2003; 83:835-870.

Witte MB, Barbul A. General principles of wound healing. Surg Clin North Am 1997; 77(3):509-528.

Xia YP, Zhao Y, Marcus J, Jiminez PA, Ruben SM, Moore PA, et al. Effects of keratinocyte growth factor-2 (KGF-2) on wound healing in an ischaemia-impaired rabbit ear model and on scar formation. J Pathol 1999; 188(4):431-438.

Author's note

Portions of this course were derived from the author's article: Krafts K. *Tissue Repair: the Hidden Drama*. Organogenesis 2010; 6. The Copyright Revision Act (PL 94-553), which became effective January 1, 1978, states that the copyright of a work is vested in the author from the moment of creation. Therefore, all authors who wish to publish in Cell Cycle must grant an exclusive license to Landes Bioscience. It is understood that the authors grant Landes Bioscience an exclusive license to publish the work and also grant rights of reproduction, derivation, distribution, sale and display. Landes Bioscience has granted back to the author the copyright permission to use portions of the above article in preparation of this course.

Course Exam: Wound Healing: A New Understanding of the Drama

1. Quiescent tissues include all of the following except:

- A. Liver
- B. Kidney
- C. Pancreas
- D. Heart

2. Which of the following cells plays a central role in scar formation?

- A. Eosinophil
- B. Mast cell
- C. Fibroblast
- D. Dendritic cell

3. The most important role of transforming growth factor beta (TGF- β) in wound healing is:

- A. Promotion of fibrosis
- B. Attraction of Neutrophils
- C. Activation of macrophages
- D. Promotion of angiogenesis

4. The end result of receptor mediated signal transduction is:
- A. Cell death
 - B. A change in gene expression
 - C. Influx of calcium into the cell
 - D. Neoplastic transformation
5. In which phase of the cell cycle is DNA synthesized?
- A. G1
 - B. S
 - C. G2
 - D. M
6. Which of the following is a function of the extra-cellular matrix?
- A. Imparts resilience to soft tissues and bone
 - B. Stores growth factors
 - C. Acts as a scaffold for migrating cells
 - D. Facilitates cell growth
 - E. All of the above
7. An injury to a continuously dividing tissue (such as oral mucosa) that disrupts the underlying stromal framework will result in:
- A. Regeneration
 - B. Scarring
8. Which of the following is true of first-intention wounds?
- A. Relatively larger than second-intention wounds
 - B. Healing is generally slow
 - C. Inflammation is abundant
 - D. Scarring is minimal
9. What happens first in the immediate aftermath of a wound?
- A. Neutrophils flood the damaged tissue
 - B. Exposed collagen activates coagulation
 - C. Epithelial cells begin to proliferate at the margins
 - D. Macrophages enter the damaged tissue
 - E. Fibroblasts secrete collagen

10. When does granulation tissue appear?
- A. Within the first few days after a wound occurs
 - B. At approximately 10 days after a wound occurs
 - C. Between two and three weeks after a wound occurs
 - D. When the final scar is being formed
11. A scar has more tensile strength than normal skin.
- A. True
 - B. False
12. A mature scar is composed of:
- A. Fibrin
 - B. Elastin
 - C. Collagen
 - D. Smooth muscle
13. An insufficiency of which vitamin causes the formation of unstable collagen?
- A. A
 - B. B
 - C. C
 - D. D
 - E. E
14. What is the single most important cause of delayed wound healing?
- A. Diabetes
 - B. Drug side effects
 - C. Foreign bodies
 - D. Size of the wound
 - E. Infection
15. Keloids contain an excess of:
- A. Collagen
 - B. Fibrin
 - C. Granulation tissue
 - D. Blood vessels
 - E. Inflammation